

## Computational Fluid Dynamics Modeling of Erosion of a Solid Bed by Jet Impingement

Authorship: Brigette Rosendall<sup>1</sup>, YoChan Kim<sup>2</sup>, Leonard J. Peltier<sup>3</sup>, Chris Kennedy<sup>4</sup>, Kelly J. Knight<sup>5</sup>, Jon Berkoe<sup>6</sup>, and Milorad Dordivich<sup>7</sup>

Bechtel National, Inc., Advanced Simulations and Analysis; <sup>1</sup>[brosenda@bechtel.com](mailto:brosenda@bechtel.com), <sup>2</sup>[ykim2@bechtel.com](mailto:ykim2@bechtel.com), <sup>3</sup>[ljpeltier@bechtel.com](mailto:ljpeltier@bechtel.com), <sup>4</sup>[ckenned@bechtel.com](mailto:ckenned@bechtel.com), <sup>5</sup>[kknight@bechtel.com](mailto:kknight@bechtel.com), <sup>6</sup>[jberkoe@bechtel.com](mailto:jberkoe@bechtel.com), <sup>7</sup>[mcdordev@bechtel.com](mailto:mcdordev@bechtel.com)

### Abstract

Computational fluid dynamics (CFD) is an emerging tool for predicting flow fields in a mixing environment. Confidence in models for solid-liquid mixing is of growing demand. Many studies have been done to look at impeller mixed systems, both in single and multi-phase systems. This work focuses on the physics of jet-induced mixing. How well can CFD predict the flow fields in single phase systems? If well enough, can we extend our confidence to solid-liquid systems?

Results of CFD models are compared to experimental data and theoretical applications from literature. First, CFD results are compared to single-phase impinging jet correlations from literature (Poreh, 1967) for wall shear and velocity decay. Next, a model of a transient, intermittent jet is compared to single-phase experimental velocity measurements made in a mixing vessel. Finally, a CFD model of a steady, clear impinging jet scouring a layer of solids is compared to data for rate of erosion and zone of influence.

**keywords:** Computational Fluid Dynamics, solid-liquid mixing, experimental validation